

University of Dundee

## Small airway dysfunction is associated with poorer asthma control

Manoharan, Arvind; Anderson, William J.; Lipworth, Joseph; Ibrahim, Ibrahim; Lipworth, Brian J.

*Published in:*  
European Respiratory Journal

*DOI:*  
[10.1183/09031936.00082314](https://doi.org/10.1183/09031936.00082314)

*Publication date:*  
2014

*Document Version*  
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

### *Citation for published version (APA):*

Manoharan, A., Anderson, W. J., Lipworth, J., Ibrahim, I., & Lipworth, B. J. (2014). Small airway dysfunction is associated with poorer asthma control. *European Respiratory Journal*, 44(5), 1353-1355.  
<https://doi.org/10.1183/09031936.00082314>

### **General rights**

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



## Small Airways Dysfunction is Associated with Poorer Asthma Control

Journal:	<i>European Respiratory Journal</i>
Manuscript ID:	ERJ-00823-2014
Manuscript Type:	Letter
Date Submitted by the Author:	04-May-2014
Complete List of Authors:	Manoharan, Arvind; University of Dundee, Scottish Centre for Respiratory Research Anderson, William; University of Dundee, Scottish Centre for Respiratory Research Lipworth, Joseph; University of Dundee, Scottish Centre for Respiratory Research Ibrahim, Ibrahim; University of Dundee, Scottish Centre for Respiratory Research Lipworth, Brian; University of Dundee, Scottish Centre for Respiratory Research
Key Words:	asthma, asthma clinical care, asthma - small airways

SCHOLARONE™  
Manuscripts

This is the peer reviewed version of the following article: Manoharan, A., et al. (2018) "Small airway dysfunction is associated with poorer asthma control", *European Respiratory Journal* 44:5, which has been published in final form at <http://dx.doi.org/10.1183/09031936.00082314>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

**Title:** Small Airways Dysfunction is Associated with Poorer Asthma Control

**Authors:** Arvind Manoharan, MBChB, William J. Anderson, MBChB, Joseph Lipworth, Ibrahim Ibrahim, and Brian J. Lipworth, MD

**Affiliations:** Scottish Centre for Respiratory Research, Ninewells Hospital and Medical School, University of Dundee, DD1 9SY, Scotland, United Kingdom

**Corresponding author:** Brian J. Lipworth, Scottish Centre for Respiratory Research, Division of Cardiovascular and Diabetes Medicine, Medical Research Institute, University of Dundee  
Ninewells Hospital and Medical School, Dundee DD1 9SY, Scotland, United Kingdom

Tel: +44 (0) 1382 383 902, Fax: +44 (0) 1382 644972

Email for correspondence: [b.j.lipworth@dundee.ac.uk](mailto:b.j.lipworth@dundee.ac.uk)

**Word count:** 1196

**Take home message**

In asthmatics with a preserved FEV<sub>1</sub>, small airways dysfunction defined by FEF<sub>25-75</sub> and R5-R20 was associated with poorer long-term control.

### Small Airways Dysfunction is Associated with Poorer Asthma Control

The clinical relevance of the small airways in persistent asthma has been gaining greater recognition in recent years [1]. Studies have shown that a significant proportion of asthmatics on standard treatment fail to achieve satisfactory asthma control. For example, in one study of 3421 asthmatic subjects who underwent guideline driven dose titration with standard inhaled corticosteroids (ICS) / long-acting beta-agonist (LABA) combination therapy over 1 year, only 41% achieved total control of their asthma while 71% were well controlled [2].

Anderson et al [3] found a high prevalence of adult patients with persistent small airway dysfunction determined by impulse oscillometry (IOS, as R5-R20) and spirometry (as FEF<sub>25-75</sub>) across British Thoracic Society (BTS) treatment steps for asthma, many of whom had a preserved FEV<sub>1</sub>. This in turn suggests an unmet clinical need in terms of patients who may have a small airway asthma phenotype.

We therefore evaluated whether small airways dysfunction was associated with worse control in adult asthmatics with a preserved FEV<sub>1</sub> (FEV<sub>1</sub> > 80% predicted). Spirometry and IOS measurements from unselected asthmatics referred from primary care who attended for screening visits for clinical trials were linked to prescription data. The prescription data were obtained from the Tayside Health Informatics Centre which links all community dispensed prescriptions using a person's unique identifier, the Community Health Index. Spirometry and IOS measurements from asthmatics were linked to oral corticosteroid and short-acting beta-agonist (SABA) use. We evaluated if small airways dysfunction, defined as FEF<sub>25-75</sub> < 70%, or peripheral airway resistance as R5-R20 > 0.07 kPa·L<sup>-1</sup>·s was associated with increased oral corticosteroid and SABA use. Oral steroid and SABA use 1 year prior and 1 year following the index measurements were determined i.e. whether or not patients had an oral steroid prescription for an asthma exacerbation or the use of > 4 or ≤ 4 SABA inhalers.

Research ethics committee approval was obtained for all the studies the patients were being screened into and Caldicott Guardian approval was obtained to transfer the data to the Health Informatics Centre. IOS (Jaeger Masterscreen IOS, Hochberg, Germany) was performed in triplicate in accordance with manufacturer’s guidelines. A SuperSpiro spirometer (Micro Medical Ltd., Chatham, Kent, United Kingdom) was used in triplicate in accordance with European Society guidelines [4]. Logistic regression analysis was applied to calculate the odds ratios (OR) for steroid and salbutamol use in the different groups. Age, gender, ICS, LABA and leukotriene receptor antagonists (LTRA) use were all included as covariates to calculate the adjusted OR and 95% confidence interval.

302 out of 442 (68%) asthmatics had a preserved  $FEV_1 > 80\%$ , mean age: 40 years,  $FEV_1$ : 97%, median ICS dose: 800 $\mu$ g, 42% taking LABA, 22% on LTRA and 5% on theophylline. The proportion of patients at BTS treatment steps 1-4 were 6.3%, 37.7%, 27.8 % and 28.0 % respectively .The results in Table 1 showed that persistent small airways dysfunction, defined by  $FEF_{25-75}$  and R5-20, was associated with a significantly increased likelihood of having worse long-term asthma control. The risk of having poorer control was greater when measurements of  $FEF_{25-75}$  and R5-R20 were combined. However, adding in  $FEV_1/FVC$  to the model did not appreciably improve the OR compared to the combined outcome of  $FEF_{25-75}$  and R5-20 because  $FEV_1/FVC$  and  $FEF_{25-75}$  were both highly correlated ( $r = 0.82$ ,  $P < 0.001$ ).

When our analysis was corrected for factors including age, gender, ICS, LABA and LTRA, the adjusted OR for  $FEF_{25-75}$  and R5-20 were similar. R5% predicted (n=302) and resonant frequency (n=268) however, did not have a significant impact in determining asthma control. There were insufficient evaluable data to perform a meaningful analysis on reactance area (AX) (n=75). During the study period, in those with a preserved  $FEV_1$ , there were a total of 14 Emergency Department visits and 33 hospital admissions for asthma exacerbations. However, these numbers were too small to perform a meaningful analysis.

Our results are similar to those of previously reported studies in asthmatic children by Shi et al [5] who showed a significant difference between selected cohorts of controlled and uncontrolled asthmatic children for both  $FEF_{25-75}$  and  $FEV_1/FVC$ , while peripheral resistance ( $R5-R20 > 0.15 \text{ kPa}\cdot\text{L}^{-1}\cdot\text{s}$ ) and reactance area ( $AX > 0.95 \text{ kPa}\cdot\text{L}^{-1}$ ) were equally predictive for detecting control. In a prospective follow up study [6] of initially controlled asthmatic children, the same authors observed a significant difference in  $FEV_1/FVC$  but not  $FEF_{25-75}$  at baseline, but for both  $FEV_1/FVC$  and  $FEF_{25-75}$  at follow up after 3 months, comparing those who remained controlled to those who subsequently became uncontrolled. Rao et al [7] in a similar design to the present study over 2 years using electronic prescribing linkage, compared matched groups of asthmatic children who had a preserved  $FEV_1$  ( $> 80\%$ ) with an abnormal  $FEV_1/FVC$  ( $< 0.85$ ) and  $FEF_{25-75}$  ( $< 60\%$ ) versus those with normal values, showing significantly increased OR for loss of control in terms of oral steroid use, asthma exacerbations and controller use.

We elected to use cut-off thresholds for small airways measurements which provided the best compromise in terms of achieving balanced numbers of patients in each group from which to make informative comparisons. Such cut-off values for normality are always going to be rather arbitrary whether they are more or less severe in nature. We acknowledge our data has some limitations in terms of it being a retrospective type health informatics study linked to a single index measurement of pulmonary function. However, we feel that our data more closely reflects real life practice where compliance is usually poor in the community. Our unselected cohort of persistent asthmatics were referred from primary care and reflected a wide spectrum of severity across BTS treatment steps.

We feel that our study may have some important potential clinical implications. It appears that effort independent (i.e. IOS) and effort dependent measurements (i.e. spirometry) may provide distinct yet complimentary information on the small airways phenotype, as shown by

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

the higher OR for the composite of FEF<sub>25-75</sub> and R5-R20 compared to either measurement alone.

It remains unclear as to whether small airway markers may be improved by using extra fine particle inhaled therapy including currently available extra fine ICS and ICS/LABA formulations and how this relates to long-term asthma control. We also do not know if small airways dysfunction as reflected by abnormal FEF<sub>25-75</sub> or R5-20 is due to ongoing persistent inflammation or simply altered airways geometry. Several prospective randomised controlled trials have shown greater improvements in small airways outcomes in response to extra fine compared to coarse particle ICS formulations in unselected patient cohorts [8-12]. Other retrospective health informatics data comparing extra fine and coarse particle ICS formulations have revealed consistent results in terms of improved asthma control based on prescribing outcomes, but have not measured any small airway pulmonary function outcomes [13-15]. We believe the time has now come for designing prospective randomized controlled trials enrolling patients with an enriched small airways phenotype, perhaps powered on pragmatic outcomes such as the Asthma Control Questionnaire.

In conclusion, we have shown that in adult asthmatics who have a preserved FEV<sub>1</sub>, the presence of persistent small airways dysfunction was associated with poorer control, perhaps suggesting the presence of a defined small airway asthma phenotype.

**Table 1** Odds ratio (95% CI) for small airway indices in 302 patients with FEV<sub>1</sub> > 80% predicted

	Crude odds ratio	P value	Adjusted odds ratio	P value
<b>FEF<sub>25-75</sub> &lt; 70 % n=157 versus FEF<sub>25-75</sub> &gt; 70% n=145</b>				
Oral steroid use	1.67 (1.04-2.68)	0.04	1.50 (0.91-2.48)	0.11
SABA use	2.00 (1.27-3.16)	0.003	1.91 (1.19—3.07)	0.007
<b>FEV<sub>1</sub>/FVC &lt; 0.80, n=167 versus FEV<sub>1</sub>/FVC &gt; 0.80, n=135</b>				
Oral steroid use	2.06 (1.27-3.35)	0.004	1.85 (1.10-3.12)	0.02
SABA use	1.61 (1.02-2.54)	0.04	1.54 (0.95-2.51)	0.08
<b>R5-R20 &gt; 0.07 kPa·L<sup>-1</sup>·s, n=135 versus R5-R20 &lt; 0.07 kPa·L<sup>-1</sup>·s, n =167</b>				
Oral steroid use	1.99 (1.23-3.19)	0.005	1.80 (1.09-2.98)	0.02
SABA use	1.83 (1.16-2.89)	0.01	1.87 (1.15-3.01)	0.01
<b>FEF<sub>25-75</sub> &lt; 70 % &amp; R5-R20 &gt; 0.07 kPa·L<sup>-1</sup>·s; n=83 versus FEF<sub>25-75</sub> &gt; 70% &amp; R5-R20 &lt; 0.07 kPa·L<sup>-1</sup>·s; n=93</b>				
Oral steroid use	2.77 (1.48-5.18)	0.001	2.34 (1.20-4.58)	0.01
SABA use	3.07 (1.66-5.67)	<0.001	3.16 (1.64-6.07)	0.001
<b>FEF<sub>25-75</sub> &lt; 70 % , R5-R20 &gt;0.07 kPa·L<sup>-1</sup>·s &amp; FEV<sub>1</sub>/FVC &lt; 0.80; n=72 versus FEF<sub>25-75</sub> &gt; 70% &amp; R5-R20 &lt; 0.07 kPa·L<sup>-1</sup>·s &amp; FEV<sub>1</sub>/FVC &gt; 0.80; n=75</b>				
Oral steroid use	3.29 (1.64-6.61)	0.001	2.78 (1.28-6.04)	0.01
SABA use	3.16 (1.61-6.19)	0.001	2.96 (1.44-6.12)	0.003



References

1. Lipworth B. Targeting the small airways asthma phenotype: if we can reach it, should we treat it? *Ann Allergy Asthma Immunol* 2013;110:233-9.

2. Bateman ED, Boushey HA, Bousquet J, et al. Can guideline-defined asthma control be achieved? The Gaining Optimal Asthma Control study. *Am J Respir Crit Care Med* 2004;170:836-44.

3. Anderson WJ, Zajda E, Lipworth BJ. Are we overlooking persistent small airways dysfunction in community-managed asthma? *Ann Allergy Asthma Immunol* 2012;109:185-9.

4. Miller MR, Hankinson J, Brusasco V, et al. Standardisation of spirometry. *Eur Respir J* 2005;26:319-38.

5. Shi Y, Aledia AS, Tatavoosian AV, et al. Relating small airways to asthma control by using impulse oscillometry in children. *J Allergy Clin Immunol* 2012;129:671-8.

6. Shi Y, Aledia AS, Galant SP, et al. Peripheral airway impairment measured by oscillometry predicts loss of asthma control in children. *J Allergy Clin Immunol* 2013;131:718-29.

7. Yamaguchi M, Niimi A, Ueda T, et al. Effect of inhaled corticosteroids on small airways in asthma: investigation using impulse oscillometry. *Pulm Pharmacol Ther* 2012;49:586-92.

8. Rao DR, Gaffin JM, Baxi SN, et al. The utility of forced expiratory flow between 25% and 75% of vital capacity in predicting childhood asthma morbidity and severity. *J Asthma* 2009;22:326-32.

9. Busse WW, Brazinsky S, Jacobson K, et al. Efficacy response of inhaled beclomethasone dipropionate in asthma is proportional to dose and is improved by formulation with a new propellant. *J Allergy Clin Immunol* 1999;104:1215-22.

- 1  
2  
3 9. Goldin JG, Tashkin DP, Kleerup EC, et al. Comparative effects of hydrofluoroalkane  
4 and chlorofluorocarbon beclomethasone dipropionate inhalation on small airways:  
5 assessment with functional helical thin-section computed tomography. *J Allergy Clin*  
6  
7  
8  
9  
10 Immunol 1999;104:S258-67.
- 11  
12 10. Nicolini G, Chetta A, Simonazzi A, et al. Both bronchial and alveolar exhaled nitric  
13 oxide are reduced with extrafine beclomethasone dipropionate in asthma. *Allergy Asthma*  
14  
15  
16  
17 Proc 2010;31:85-90.
- 18  
19  
20 11. Cohen J, Postma DS, Douma WR, et al. Particle size matters: diagnostics and  
21 treatment of small airways involvement in asthma. *Eur Respir J* 2011;37:532-40.
- 22  
23  
24 12. Juniper EF, Price DB, Stampone PA, et al. Clinically important improvements in  
25 asthma-specific quality of life, but no difference in conventional clinical indexes in patients  
26 changed from conventional beclomethasone dipropionate to approximately half the dose of  
27 extrafine beclomethasone dipropionate. *Chest* 2002;121:1824-32.3.
- 28  
29  
30  
31  
32 13. Price D, Martin RJ, Barnes N, Dorinsky P, Israel E, Roche N, et al. Prescribing  
33 practices and asthma control with hydrofluoroalkane-beclomethasone and fluticasone: a real-  
34 world observational study. *J Allergy Clin Immunol* 2010;126:511-8 e1-10.
- 35  
36  
37  
38 14. Barnes N, Price D, Colice G, et al. Asthma control with extrafine-particle  
39 hydrofluoroalkane-beclometasone vs. large-particle chlorofluorocarbon-beclometasone: a  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60 real-world observational study. *Clin Exp Allergy* 2011;41:1521-32.
15. Colice G, Martin RJ, Israel E, et al. Asthma outcomes and costs of therapy with  
extrafine beclomethasone and fluticasone. *J Allergy Clin Immunol* 2013;132:45-54.